

Colloquium

Numerical evidence for the L^p -norm blow-up

主講人：卓建宏 教授
國立中正大學數學系

時 間：106 年 11 月 1 日(三) 14 : 30

地 點：應用數學系多媒體教室（理 408 室）

摘 要：

We consider those evolution equations whose solutions blow up in finite time T in the sense that $\|u(t, \cdot)\|_\infty \rightarrow \infty$ as $t \rightarrow T$. Here, $u(t, x)$ denotes the exact solution of the evolution equation under consideration. Such problems are often called blow-up problems and the finite time is called the blow-up time. Although the L^∞ -norm for the solutions of such problems becomes unbounded in a finite time T , the L^p -norms ($1 \leq p < \infty$) do not always blow up simultaneously. However, for the numerical computation of blow-up problems, if we consider time-independent spatial partitions for a bounded domain to compute blow-up solutions, the unboundedness for the discrete L^∞ -norm of the numerical solution is equivalent to the unboundedness of any other discrete L^p -norms ($1 \leq p < \infty$) for the numerical solutions. That is, all discrete L^p -norms blow up simultaneously while the L^p -norms of the exact solution may remain bounded as $t \rightarrow T$. This shows the numerical difficulty in the computation of L^p -norms of the blow-up solutions which remain bounded as $t \rightarrow T$. We would like to explore in this talk such a problem by a simple model equation axisymmetric semi-linear heat equation. Some remarks on the numerical blow-up solutions of axisymmetric semi-linear equation will also be reported.